

**Table 10.6d**  
**Immunosuppression Use for Maintenance by Regimen**  
**Prior to Discharge, 1999 to 2008**  
**Recipients with Intestine Transplants**

	Transplant Year									
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
<b>Functioning Graft at Discharge</b>	49	61	88	89	103	130	155	148	166	168
<b>With Immunosuppression Info</b>	49	61	88	89	103	127	152	148	166	167
Tac	4.1%	1.6%	18.2%	31.5%	30.1%	22.8%	25.7%	16.9%	11.4%	10.2%
--- + MMF/MPA	0.0%	1.6%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + MMF/MPA	2.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.3%	7.4%	11.4%	10.8%
Tac + Siro/Evero	0.0%	0.0%	2.3%	0.0%	0.0%	2.4%	2.0%	2.0%	0.0%	0.6%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	1.0%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + Steroids	0.0%	0.0%	1.1%	2.2%	3.9%	0.8%	2.0%	1.4%	1.8%	0.0%
CyA + Steroids	4.1%	0.0%	0.0%	0.0%	1.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + Steroids	67.3%	41.0%	48.9%	36.0%	50.5%	55.9%	47.4%	52.7%	56.0%	55.1%
--- + MMF/MPA + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	0.7%	0.0%	0.6%
Tac + MMF/MPA + Steroids	14.3%	13.1%	3.4%	7.9%	6.8%	8.7%	10.5%	10.1%	8.4%	12.6%
--- + OtherAntimet + Steroids	0.0%	1.6%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + OtherAntimet + Steroids	6.1%	0.0%	6.8%	0.0%	0.0%	0.0%	0.0%	0.7%	1.8%	0.0%
--- + Siro/Evero + Steroids	0.0%	1.6%	1.1%	0.0%	0.0%	1.6%	0.7%	0.0%	0.0%	0.0%
Tac + Siro/Evero + Steroids	0.0%	34.4%	18.2%	15.7%	3.9%	7.9%	9.9%	7.4%	7.8%	9.0%
Other Regimen	2.0%	4.9%	0.0%	6.7%	2.9%	0.0%	0.0%	0.7%	1.2%	1.2%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Percentages in each column sum to 100 percent.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.  
 "Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.