

Table 10.6e
Immunosuppression Use for Maintenance
Prior to Discharge, 1999 to 2008
Recipients with Intestine Transplants

	Year of Transplant									
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Functioning Graft at Discharge	49	61	88	89	103	130	155	148	166	168
With Immunosuppression Info	49	61	88	89	103	127	152	148	166	167
Corticosteroids										
Any in Category	93.9%	96.7%	79.5%	66.3%	67.0%	74.8%	71.1%	73.6%	77.1%	78.4%
Steroids	93.9%	96.7%	79.5%	66.3%	67.0%	74.8%	71.1%	73.6%	77.1%	78.4%
Cyclosporine										
Any in Category	4.1%	0.0%	0.0%	0.0%	1.0%	0.0%	0.0%	0.0%	0.0%	0.6%
Neoral	4.1%	0.0%	0.0%	0.0%	1.0%	0.0%	0.0%	0.0%	0.0%	0.6%
Tacrolimus										
Any in Category	95.9%	95.1%	97.7%	95.5%	93.2%	97.6%	96.7%	98.0%	98.2%	99.4%
Tacrolimus	95.9%	95.1%	97.7%	95.5%	93.2%	97.6%	96.7%	98.0%	97.0%	98.8%
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.2%	0.6%
Antimetabolites										
Any in Category	24.5%	21.3%	10.2%	12.4%	8.7%	8.7%	12.5%	19.6%	22.9%	24.6%
Mycophenolate Mofetil	18.4%	14.8%	3.4%	11.2%	8.7%	8.7%	12.5%	18.9%	21.1%	24.0%
Mycophenolate Sodium	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Azathioprine	6.1%	6.6%	6.8%	1.1%	0.0%	0.0%	0.0%	0.7%	1.8%	0.6%
mTOR Inhibitors										
Any in Category	2.0%	39.3%	21.6%	20.2%	5.8%	11.8%	12.5%	10.1%	9.0%	9.6%
Sirolimus	2.0%	39.3%	21.6%	20.2%	5.8%	11.8%	12.5%	10.1%	9.0%	9.6%
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Individual drug percentages will not necessarily add up to the drug category percentages, as patients may be prescribed more than one drug within the same category.