

Table 10.8
Adjusted Graft Survival, Intestine Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|---------------------------------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2006 - 2007) | | | (Tx 2006 - 2007) | | | (Tx 2002 - 2007) | | | (Tx 1997 - 2007) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 342 | 86.9% | 1.8% | 342 | 73.2% | 2.4% | 860 | 48.4% | 2.4% | 1,242 | 34.4% | 2.4% |
| Age at Tx | | | | | | | | | | | | |
| <1 Year | 64 | 79.5% | 5.0% | 64 | 63.6% | 6.0% | 133 | 39.8% | 6.1% | 188 | 30.2% | 5.6% |
| 1-5 Years | 101 | 85.1% | 3.5% | 101 | 72.3% | 4.4% | 266 | 52.4% | 4.1% | 403 | 41.1% | 3.6% |
| 6-11 Years | 14 | 76.9% | 11.2% | 14 | 61.5% | 12.8% | 57 | 38.4% | 8.4% | 91 | 37.1% | 7.2% |
| 12-17 Years | 16 | 87.5% | 8.0% | 16 | 75.0% | 10.4% | 48 | 51.4% | 8.5% | 66 | 40.2% | 8.8% |
| 18-34 Years | 57 | 87.7% | 4.3% | 57 | 73.6% | 5.8% | 128 | 59.0% | 5.3% | 176 | 36.8% | 6.2% |
| 35-49 Years | 54 | 90.7% | 3.9% | 54 | 75.9% | 5.8% | 137 | 52.5% | 6.1% | 203 | 25.6% | 6.9% |
| 50-64 Years | 32 | 96.9% | 3.0% | 32 | 84.4% | 6.3% | 84 | 29.7% | 9.2% | 107 | 12.6% | 7.1% |
| 65+ Years | 4 | + | + | 4 | + | + | 7 | 0.0% | 0.0% | 8 | 37.5% | 20.1% |
| Recipient Ethnicity/Race | | | | | | | | | | | | |
| White | 236 | 87.9% | 2.1% | 236 | 74.4% | 2.9% | 601 | 47.3% | 3.0% | 860 | 35.0% | 2.7% |
| African American | 51 | 91.1% | 3.8% | 51 | 74.4% | 6.0% | 116 | 46.2% | 6.1% | 177 | 25.4% | 5.9% |
| Hispanic/Latino | 43 | 78.8% | 6.0% | 43 | 65.8% | 7.1% | 116 | 57.3% | 5.3% | 171 | 38.2% | 7.8% |
| Asian | 6 | 69.7% | 16.2% | 6 | 69.7% | 16.2% | 16 | 49.6% | 12.2% | 22 | 44.8% | 10.4% |
| Other/Multi-race | 6 | 85.5% | 12.3% | 6 | 70.5% | 16.0% | 11 | 28.1% | 15.4% | 12 | 38.7% | 15.7% |
| Recipient Gender | | | | | | | | | | | | |
| Female | 167 | 89.4% | 2.4% | 167 | 76.8% | 3.3% | 432 | 47.7% | 3.6% | 604 | 33.4% | 3.7% |
| Male | 175 | 84.5% | 2.7% | 175 | 69.7% | 3.5% | 428 | 49.0% | 3.3% | 638 | 35.4% | 3.0% |

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on intestine transplants.

Multi-organ transplants other than intestine with liver and/or pancreas are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 10.8 (Continued)
Adjusted Graft Survival, Intestine Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|--------------------------|------------------|--------|-----------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2006 - 2007) | | | (Tx 2006 - 2007) | | | (Tx 2002 - 2007) | | | (Tx 1997 - 2007) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 342 | 86.9% | 1.8% | 342 | 73.2% | 2.4% | 860 | 48.4% | 2.4% | 1,242 | 34.4% | 2.4% |
| Primary Diagnosis | | | | | | | | | | | | |
| Short Gut Syndrome | 246 | 85.9% | 2.2% | 246 | 71.9% | 2.9% | 622 | 48.7% | 2.7% | 915 | 33.7% | 2.8% |
| Functional Bowel Problem | 48 | 84.9% | 4.9% | 48 | 73.4% | 6.2% | 149 | 43.2% | 6.4% | 214 | 34.1% | 5.0% |
| Other | 46 | 95.2% | 3.3% | 46 | 85.7% | 5.4% | 86 | 57.8% | 7.7% | 109 | 42.3% | 8.3% |
| Unknown | 2 | 100.0% | 0.0% | 2 | 0.0% | 0.0% | 3 | 0.0% | 0.0% | 4 | 0.0% | 0.0% |

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on intestine transplants.

Multi-organ transplants other than intestine with liver and/or pancreas are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 10.8 (Continued)
Adjusted Graft Survival, Intestine Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|--------------------------------|------------------|--------|-----------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2006 - 2007) | | | (Tx 2006 - 2007) | | | (Tx 2002 - 2007) | | | (Tx 1997 - 2007) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 342 | 86.9% | 1.8% | 342 | 73.2% | 2.4% | 860 | 48.4% | 2.4% | 1,242 | 34.4% | 2.4% |
| Transplant Center State | | | | | | | | | | | | |
| Alabama | 0 | - | - | 0 | - | - | 0 | - | - | 1 | 0.0% | 0.0% |
| California | 32 | 73.2% | 7.7% | 32 | 64.1% | 8.4% | 92 | 59.6% | 5.5% | 112 | 46.5% | 8.0% |
| District of Columbia | 39 | 87.4% | 5.2% | 39 | 74.5% | 6.9% | 64 | 65.3% | 7.2% | 64 | 65.1% | 7.2% |
| Florida | 49 | 78.6% | 5.7% | 49 | 64.8% | 6.7% | 136 | 35.7% | 5.3% | 238 | 29.1% | 3.8% |
| Illinois | 5 | 61.6% | 18.8% | 5 | 61.6% | 18.8% | 15 | 60.4% | 12.0% | 20 | 42.9% | 11.1% |
| Indiana | 28 | 88.8% | 6.0% | 28 | 73.5% | 8.5% | 49 | 57.5% | 8.1% | 49 | 57.8% | 8.1% |
| Louisiana | 0 | - | - | 0 | - | - | 1 | 0.0% | 0.0% | 1 | 0.0% | 0.0% |
| Massachusetts | 6 | 84.1% | 13.3% | 6 | 68.1% | 16.9% | 8 | 74.7% | 14.4% | 8 | 74.6% | 14.5% |
| Maryland | 0 | - | - | 0 | - | - | 0 | - | - | 1 | 0.0% | 0.0% |
| Minnesota | 3 | 35.5% | 20.3% | 3 | 35.5% | 20.3% | 4 | 24.1% | 16.3% | 11 | 0.0% | 0.0% |
| Missouri | 0 | - | - | 0 | - | - | 0 | - | - | 1 | 0.0% | 0.0% |
| Nebraska | 53 | 96.3% | 2.6% | 53 | 77.3% | 5.8% | 131 | 45.3% | 7.9% | 219 | 39.3% | 4.6% |
| New York | 14 | 85.0% | 9.5% | 14 | 55.4% | 13.0% | 49 | 35.5% | 7.6% | 99 | 32.7% | 5.7% |
| Ohio | 12 | 76.2% | 11.6% | 12 | 60.1% | 13.5% | 23 | 28.5% | 11.5% | 23 | 27.7% | 11.4% |
| Oklahoma | 0 | - | - | 0 | - | - | 1 | 0.0% | 0.0% | 1 | 0.0% | 0.0% |
| Pennsylvania | 90 | 97.7% | 1.6% | 90 | 85.9% | 3.8% | 264 | 54.7% | 4.4% | 359 | 34.5% | 5.0% |
| South Carolina | 1 | 100.0% | 0.0% | 1 | 0.0% | 0.0% | 4 | 29.2% | 17.7% | 8 | 13.5% | 9.3% |
| Texas | 6 | 52.0% | 18.0% | 6 | 52.0% | 18.0% | 7 | 56.6% | 17.1% | 9 | 54.9% | 15.5% |
| Virginia | 1 | + | + | 1 | + | + | 1 | + | + | 1 | + | + |
| Washington | 3 | 68.2% | 21.6% | 3 | 68.2% | 21.6% | 3 | 66.8% | 22.2% | 3 | 66.2% | 22.4% |
| Wisconsin | 0 | - | - | 0 | - | - | 8 | 25.2% | 13.3% | 14 | 22.3% | 11.1% |

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on intestine transplants.

Multi-organ transplants other than intestine with liver and/or pancreas are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.