

Table 5.9b
Adjusted Graft Survival by Year of Transplant at 3 Months, 1 Year,
3 Years, 5 Years and 10 Years
Deceased Donor ECD Kidney Transplants

Trans- # plant Year	Trans- plants	3 Months		1 Year		3 Years		5 Years		10 Years	
		Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.
1987	43	76.0%	6.5%	66.3%	7.2%	52.6%	7.8%	36.2%	7.6%	19.4%	6.3%
1988	177	70.9%	3.5%	60.3%	3.7%	42.4%	3.8%	28.9%	3.5%	16.0%	2.8%
1989	220	80.2%	2.7%	65.6%	3.3%	48.2%	3.4%	35.4%	3.3%	14.2%	2.4%
1990	334	80.6%	2.2%	66.8%	2.6%	50.3%	2.8%	41.3%	2.8%	17.7%	2.1%
1991	396	84.5%	1.8%	75.3%	2.2%	58.7%	2.5%	42.7%	2.5%	15.9%	1.9%
1992	485	82.8%	1.7%	70.4%	2.1%	57.1%	2.3%	43.2%	2.3%	17.7%	1.8%
1993	520	81.4%	1.7%	70.7%	2.0%	54.5%	2.2%	40.7%	2.2%	17.6%	1.7%
1994	861	86.6%	1.2%	77.7%	1.4%	61.1%	1.7%	45.7%	1.7%	20.9%	1.4%
1995	1,004	87.2%	1.1%	77.4%	1.3%	63.4%	1.5%	48.6%	1.6%	24.2%	1.4%
1996	1,074	86.8%	1.0%	79.9%	1.2%	67.3%	1.4%	51.3%	1.5%	25.9%	1.4%
1997	1,131	89.5%	0.9%	81.4%	1.2%	65.3%	1.4%	51.2%	1.5%	24.8%	1.3%
1998	1,213	89.8%	0.9%	80.6%	1.1%	64.8%	1.4%	52.0%	1.4%	27.0%	1.3%
1999	1,226	89.6%	0.9%	81.6%	1.1%	67.7%	1.3%	53.6%	1.4%	+	+
2000	1,170	90.0%	0.9%	80.4%	1.2%	65.7%	1.4%	52.1%	1.5%	+	+
2001	1,188	89.9%	0.9%	80.6%	1.1%	68.5%	1.3%	54.0%	1.5%	+	+
2002	1,270	88.8%	0.9%	80.8%	1.1%	67.5%	1.3%	55.0%	1.4%	+	+
2003	1,442	90.9%	0.7%	82.6%	1.0%	68.9%	1.2%	57.6%	1.3%	+	+
2004	1,524	92.1%	0.7%	84.0%	0.9%	69.9%	1.2%	+	+	+	+
2005	1,727	92.8%	0.6%	84.7%	0.8%	73.2%	1.1%	+	+	+	+
2006	1,756	92.2%	0.6%	85.5%	0.8%	+	+	+	+	+	+
2007	1,791	92.9%	0.6%	86.7%	0.8%	+	+	+	+	+	+

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up.

Multi-organ transplants are excluded for kidney, pancreas, heart, lung, and liver survival rates.

Intestine survival includes intestine only and intestine with liver and/or pancreas transplants.

Living donor transplants excluded unless explicitly listed. Heterotopic heart and liver transplants excluded.

Patient survival follows patients from first transplant of this type until death. Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type organ.

Survival rates are adjusted to the characteristics of transplants in year 1998. See Technical Notes for details of patient and graft survival methods.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.