

Table 7.12
Adjusted Patient Survival, Pancreas After Kidney (PAK) Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	404	98.8%	0.5%	404	97.1%	0.8%	1,578	84.3%	1.2%	2,356	67.6%	2.0%
Age at Tx												
6-11 Years	0	-	-	0	-	-	1	+	+	1	+	+
18-34 Years	59	98.3%	1.7%	59	98.3%	1.7%	285	87.7%	2.7%	435	69.1%	4.8%
35-49 Years	245	99.2%	0.6%	245	97.1%	1.1%	986	84.8%	1.5%	1,501	68.9%	2.4%
50-64 Years	99	98.0%	1.4%	99	96.0%	2.0%	305	78.7%	3.3%	417	61.7%	4.4%
65+ Years	1	+	+	1	+	+	1	+	+	2	+	+
Recipient Ethnicity/Race												
White	352	98.9%	0.5%	352	97.6%	0.8%	1,345	84.1%	1.3%	2,036	67.5%	2.1%
African American	29	96.3%	3.6%	29	92.5%	5.0%	120	81.1%	4.3%	171	56.2%	8.6%
Hispanic/Latino	21	100.0%	0.0%	21	95.2%	4.6%	95	89.1%	4.7%	123	88.1%	4.0%
Asian	1	+	+	1	+	+	10	90.0%	9.0%	13	92.3%	7.1%
Other/Multi-race	1	+	+	1	+	+	8	+	+	13	78.8%	12.6%
Recipient Gender												
Female	172	98.2%	1.0%	172	97.1%	1.3%	660	84.7%	1.9%	982	68.5%	3.1%
Male	232	99.2%	0.6%	232	97.1%	1.1%	918	84.1%	1.6%	1,374	66.9%	2.6%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on pancreas and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 7.12 (Continued)
Adjusted Patient Survival, Pancreas After Kidney (PAK) Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	404	98.8%	0.5%	404	97.1%	0.8%	1,578	84.3%	1.2%	2,356	67.6%	2.0%
Transplant Center State												
Alabama	7	+	+	7	+	+	26	95.3%	4.5%	26	95.3%	4.5%
Arkansas	0	-	-	0	-	-	2	51.4%	24.7%	3	67.3%	21.9%
Arizona	16	+	+	16	+	+	50	84.8%	7.4%	55	85.5%	6.4%
California	23	+	+	23	+	+	94	85.2%	4.6%	123	86.8%	3.5%
Colorado	2	+	+	2	+	+	8	+	+	10	79.8%	16.1%
Connecticut	2	+	+	2	+	+	14	89.0%	9.8%	39	57.6%	10.1%
District of Columbia	2	50.3%	25.9%	2	50.3%	25.9%	19	94.8%	5.0%	32	54.1%	14.0%
Florida	2	+	+	2	+	+	26	77.8%	8.8%	50	71.7%	7.0%
Georgia	3	+	+	3	+	+	15	84.9%	9.6%	18	81.0%	9.6%
Hawaii	0	-	-	0	-	-	1	+	+	2	+	+
Iowa	1	+	+	1	+	+	4	+	+	7	30.3%	18.2%
Illinois	11	+	+	11	+	+	47	86.6%	5.7%	64	50.8%	12.3%
Indiana	14	+	+	14	+	+	51	95.6%	3.0%	54	0.0%	0.0%
Kansas	0	-	-	0	-	-	7	+	+	11	91.0%	8.2%
Kentucky	3	+	+	3	+	+	6	+	+	10	+	+
Louisiana	2	+	+	2	+	+	20	76.0%	10.3%	30	66.7%	10.8%
Massachusetts	21	+	+	21	+	+	86	90.6%	3.8%	119	76.4%	6.6%
Maryland	15	+	+	15	+	+	65	87.8%	4.3%	142	60.3%	6.9%
Maine	5	+	+	5	+	+	16	65.6%	17.3%	16	65.3%	17.4%
Michigan	18	87.5%	8.2%	18	87.5%	8.2%	52	66.3%	10.2%	81	65.2%	8.0%
Minnesota	41	100.0%	0.0%	41	97.7%	2.2%	209	82.9%	3.3%	467	69.0%	3.3%
Missouri	7	+	+	7	+	+	12	90.9%	8.3%	12	90.9%	8.3%
North Carolina	4	+	+	4	+	+	46	89.5%	5.1%	59	77.6%	11.6%
North Dakota	3	+	+	3	+	+	27	93.9%	5.8%	29	94.5%	5.2%
Nebraska	11	+	+	11	+	+	24	+	+	36	71.1%	18.2%
New Hampshire	9	100.0%	0.0%	9	89.7%	9.3%	11	90.9%	8.3%	11	90.9%	8.3%
New Jersey	11	100.0%	0.0%	11	92.0%	7.5%	64	75.8%	6.4%	68	68.6%	7.9%
Nevada	0	-	-	0	-	-	0	-	-	1	+	+
New York	25	+	+	25	+	+	75	79.7%	7.0%	100	51.9%	14.2%
Ohio	40	+	+	40	+	+	112	69.7%	7.0%	159	67.8%	5.5%
Oklahoma	2	+	+	2	+	+	8	+	+	8	+	+
Oregon	5	+	+	5	+	+	13	78.7%	13.1%	15	54.1%	16.8%
Pennsylvania	21	+	+	21	+	+	92	89.2%	3.4%	120	78.7%	5.8%

(Continued)

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on pancreas and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 7.12 (Continued)
Adjusted Patient Survival, Pancreas After Kidney (PAK) Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Transplant Center State												
Rhode Island	5	+	+	5	+	+	20	84.9%	9.5%	20	84.9%	9.5%
South Carolina	5	100.0%	0.0%	5	81.6%	14.8%	14	62.6%	15.2%	18	0.0%	0.0%
South Dakota	8	100.0%	0.0%	8	88.4%	10.4%	22	82.8%	9.0%	22	82.7%	9.0%
Tennessee	6	+	+	6	+	+	32	96.9%	3.0%	45	72.6%	11.3%
Texas	20	100.0%	0.0%	20	95.4%	4.5%	53	76.6%	10.4%	84	50.9%	11.1%
Utah	3	+	+	3	+	+	24	91.0%	6.0%	28	92.4%	5.1%
Virginia	13	84.6%	9.6%	13	84.6%	9.6%	47	95.8%	2.9%	53	96.2%	2.6%
Vermont	1	+	+	1	+	+	3	+	+	3	+	+
Washington	5	+	+	5	+	+	12	68.2%	18.5%	25	71.6%	9.8%
Wisconsin	12	100.0%	0.0%	12	91.2%	8.0%	49	84.3%	6.1%	81	78.9%	5.8%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on pancreas and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.