

Table 8.8
Adjusted Kidney Graft Survival, Kidney-Pancreas Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|---------------------------------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2006 - 2007) | | | (Tx 2006 - 2007) | | | (Tx 2002 - 2007) | | | (Tx 1997 - 2007) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 1,761 | 96.0% | 0.5% | 1,761 | 92.6% | 0.6% | 5,304 | 78.7% | 0.8% | 9,836 | 58.7% | 0.9% |
| Age at Tx | | | | | | | | | | | | |
| 1-5 Years | 0 | - | - | 0 | - | - | 0 | - | - | 1 | + | + |
| 6-11 Years | 0 | - | - | 0 | - | - | 1 | + | + | 1 | + | + |
| 12-17 Years | 0 | - | - | 0 | - | - | 1 | + | + | 2 | + | + |
| 18-34 Years | 351 | 96.2% | 1.0% | 351 | 91.2% | 1.5% | 1,211 | 74.1% | 1.7% | 2,529 | 53.2% | 1.7% |
| 35-49 Years | 1,061 | 96.1% | 0.6% | 1,061 | 93.1% | 0.8% | 3,171 | 80.6% | 1.0% | 5,887 | 60.8% | 1.1% |
| 50-64 Years | 344 | 95.4% | 1.1% | 344 | 92.1% | 1.5% | 909 | 76.9% | 2.0% | 1,403 | 58.0% | 2.9% |
| 65+ Years | 5 | + | + | 5 | + | + | 11 | 90.9% | 8.3% | 13 | 58.9% | 17.7% |
| Recipient Ethnicity/Race | | | | | | | | | | | | |
| White | 1,306 | 96.4% | 0.5% | 1,306 | 93.2% | 0.7% | 3,961 | 80.4% | 0.9% | 7,710 | 59.6% | 1.0% |
| African American | 265 | 94.5% | 1.4% | 265 | 89.6% | 1.9% | 780 | 68.5% | 2.6% | 1,256 | 51.0% | 2.8% |
| Hispanic/Latino | 148 | 94.7% | 1.8% | 148 | 91.3% | 2.3% | 452 | 78.6% | 2.6% | 706 | 61.7% | 3.6% |
| Asian | 23 | + | + | 23 | + | + | 75 | 87.9% | 4.5% | 108 | 63.7% | 9.2% |
| Other/Multi-race | 19 | 94.8% | 4.9% | 19 | 89.6% | 6.8% | 36 | 66.5% | 9.8% | 56 | 40.3% | 12.3% |
| Recipient Gender | | | | | | | | | | | | |
| Female | 650 | 96.4% | 0.7% | 650 | 93.4% | 1.0% | 2,012 | 77.0% | 1.3% | 3,899 | 56.5% | 1.5% |
| Male | 1,111 | 95.7% | 0.6% | 1,111 | 92.0% | 0.8% | 3,292 | 79.7% | 1.0% | 5,937 | 60.1% | 1.1% |

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney-pancreas and kidney transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 8.8 (Continued)
Adjusted Kidney Graft Survival, Kidney-Pancreas Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|--------------------------------|------------------|--------|-----------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2006 - 2007) | | | (Tx 2006 - 2007) | | | (Tx 2002 - 2007) | | | (Tx 1997 - 2007) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 1,761 | 96.0% | 0.5% | 1,761 | 92.6% | 0.6% | 5,304 | 78.7% | 0.8% | 9,836 | 58.7% | 0.9% |
| Transplant Center State | | | | | | | | | | | | |
| Alabama | 37 | + | + | 37 | + | + | 97 | 79.8% | 5.5% | 167 | 65.5% | 5.9% |
| Arkansas | 5 | + | + | 5 | + | + | 18 | 87.0% | 8.4% | 46 | 59.8% | 10.3% |
| Arizona | 70 | 98.6% | 1.4% | 70 | 95.8% | 2.3% | 132 | 83.8% | 5.3% | 182 | 66.3% | 6.3% |
| California | 214 | 95.8% | 1.4% | 214 | 90.9% | 2.0% | 660 | 81.4% | 2.0% | 1,199 | 59.3% | 3.0% |
| Colorado | 25 | 96.0% | 3.9% | 25 | 96.0% | 3.9% | 75 | 90.3% | 5.0% | 154 | 59.9% | 7.5% |
| Connecticut | 2 | + | + | 2 | + | + | 9 | 89.4% | 9.4% | 25 | 60.7% | 11.1% |
| District of Columbia | 13 | 92.6% | 6.9% | 13 | 85.1% | 9.3% | 51 | 74.2% | 7.8% | 119 | 47.0% | 9.1% |
| Florida | 133 | 96.2% | 1.6% | 133 | 92.4% | 2.3% | 428 | 78.9% | 2.8% | 648 | 54.9% | 4.7% |
| Georgia | 48 | 95.8% | 2.9% | 48 | 93.6% | 3.5% | 150 | 79.6% | 4.7% | 274 | 62.2% | 5.1% |
| Hawaii | 5 | + | + | 5 | + | + | 13 | 80.7% | 12.0% | 20 | 0.0% | 0.0% |
| Iowa | 13 | + | + | 13 | + | + | 53 | 91.2% | 4.3% | 118 | 77.6% | 5.1% |
| Illinois | 77 | 97.4% | 1.8% | 77 | 96.1% | 2.2% | 350 | 73.5% | 3.1% | 703 | 52.3% | 3.2% |
| Indiana | 59 | 98.3% | 1.6% | 59 | 98.3% | 1.6% | 151 | 75.4% | 6.3% | 183 | 52.0% | 13.3% |
| Kansas | 15 | 93.4% | 6.1% | 15 | 85.7% | 9.0% | 45 | 79.9% | 7.0% | 94 | 74.4% | 5.4% |
| Kentucky | 25 | 92.1% | 5.2% | 25 | 88.1% | 6.3% | 82 | 67.5% | 6.3% | 161 | 47.9% | 6.5% |
| Louisiana | 41 | 95.1% | 3.3% | 41 | 90.3% | 4.6% | 149 | 72.5% | 4.5% | 245 | 50.9% | 6.6% |
| Massachusetts | 7 | + | + | 7 | + | + | 22 | 95.5% | 4.3% | 47 | 75.5% | 8.1% |
| Maryland | 64 | 93.8% | 3.0% | 64 | 85.5% | 4.5% | 120 | 82.2% | 4.3% | 240 | 56.5% | 5.4% |
| Maine | 4 | + | + | 4 | + | + | 13 | 92.6% | 6.9% | 13 | 92.5% | 7.0% |
| Michigan | 51 | 92.2% | 3.7% | 51 | 90.3% | 4.1% | 110 | 79.3% | 6.8% | 173 | 65.8% | 5.8% |
| Minnesota | 45 | 88.6% | 4.8% | 45 | 88.6% | 4.8% | 132 | 63.2% | 5.1% | 354 | 51.1% | 3.5% |
| Missouri | 27 | 100.0% | 0.0% | 27 | 96.3% | 3.5% | 70 | 55.2% | 12.3% | 111 | 42.6% | 8.3% |
| North Carolina | 87 | 96.5% | 2.0% | 87 | 94.2% | 2.5% | 219 | 86.9% | 2.8% | 359 | 62.9% | 4.6% |
| North Dakota | 5 | + | + | 5 | + | + | 16 | 84.3% | 10.0% | 16 | 84.4% | 9.9% |
| Nebraska | 19 | + | + | 19 | + | + | 56 | 75.0% | 8.0% | 113 | 75.0% | 5.7% |
| New Hampshire | 7 | + | + | 7 | + | + | 7 | + | + | 7 | + | + |
| New Jersey | 42 | 93.0% | 3.9% | 42 | 93.0% | 3.9% | 99 | 77.1% | 5.7% | 180 | 69.0% | 5.0% |
| Nevada | 0 | - | - | 0 | - | - | 3 | + | + | 14 | + | + |
| New York | 34 | 94.3% | 3.9% | 34 | 88.6% | 5.3% | 110 | 70.4% | 6.2% | 262 | 63.9% | 4.1% |
| Ohio | 123 | 95.1% | 1.9% | 123 | 89.4% | 2.8% | 344 | 80.6% | 2.5% | 646 | 56.3% | 3.2% |
| Oklahoma | 12 | 100.0% | 0.0% | 12 | 91.1% | 8.0% | 34 | 63.9% | 11.8% | 56 | 53.7% | 10.9% |
| Oregon | 21 | 100.0% | 0.0% | 21 | 94.6% | 5.1% | 54 | 81.4% | 6.6% | 84 | 68.8% | 7.9% |
| Pennsylvania | 78 | 94.8% | 2.5% | 78 | 93.5% | 2.8% | 299 | 76.1% | 3.2% | 662 | 52.4% | 3.4% |

(Continued)

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+)= Values not determined due to insufficient follow-up. (-)= Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney-pancreas and kidney transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 8.8 (Continued)
Adjusted Kidney Graft Survival, Kidney-Pancreas Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|--------------------------------|------------------|--------|-----------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2006 - 2007) | | | (Tx 2006 - 2007) | | | (Tx 2002 - 2007) | | | (Tx 1997 - 2007) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Transplant Center State | | | | | | | | | | | | |
| Puerto Rico | 5 | 81.3% | 15.2% | 5 | 81.3% | 15.2% | 5 | 79.5% | 16.4% | 5 | 79.5% | 16.5% |
| Rhode Island | 2 | 100.0% | 0.0% | 2 | 46.3% | 24.7% | 5 | 79.6% | 16.4% | 5 | 79.9% | 16.2% |
| South Carolina | 22 | 100.0% | 0.0% | 22 | 95.2% | 4.6% | 68 | 76.4% | 7.5% | 141 | 53.2% | 6.8% |
| South Dakota | 0 | - | - | 0 | - | - | 2 | + | + | 2 | + | + |
| Tennessee | 52 | 100.0% | 0.0% | 52 | 94.2% | 3.2% | 130 | 72.5% | 5.9% | 264 | 54.3% | 4.8% |
| Texas | 86 | 96.5% | 2.0% | 86 | 93.0% | 2.8% | 285 | 85.3% | 2.6% | 519 | 67.7% | 4.2% |
| Utah | 23 | 82.4% | 7.8% | 23 | 82.4% | 7.8% | 67 | 87.0% | 4.9% | 104 | 71.4% | 5.8% |
| Virginia | 28 | 93.1% | 4.6% | 28 | 79.2% | 7.4% | 77 | 77.6% | 6.6% | 164 | 55.4% | 6.4% |
| Vermont | 5 | + | + | 5 | + | + | 14 | 73.2% | 16.2% | 15 | 0.0% | 0.0% |
| Washington | 45 | 97.8% | 2.1% | 45 | 95.6% | 3.0% | 134 | 87.2% | 4.2% | 252 | 60.9% | 5.5% |
| Wisconsin | 85 | 97.7% | 1.6% | 85 | 96.5% | 2.0% | 346 | 79.3% | 2.8% | 690 | 63.6% | 2.9% |

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney-pancreas and kidney transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 8.8 (Continued)
Adjusted Pancreas Graft Survival, Kidney-Pancreas Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|---------------------------------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2006 - 2007) | | | (Tx 2006 - 2007) | | | (Tx 2002 - 2007) | | | (Tx 1997 - 2007) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 1,761 | 88.7% | 0.8% | 1,761 | 84.8% | 0.9% | 5,304 | 73.5% | 0.8% | 9,836 | 55.3% | 0.9% |
| Age at Tx | | | | | | | | | | | | |
| 1-5 Years | 0 | - | - | 0 | - | - | 0 | - | - | 1 | + | + |
| 6-11 Years | 0 | - | - | 0 | - | - | 1 | + | + | 1 | + | + |
| 12-17 Years | 0 | - | - | 0 | - | - | 1 | + | + | 2 | + | + |
| 18-34 Years | 351 | 89.7% | 1.6% | 351 | 85.6% | 1.9% | 1,211 | 69.8% | 1.7% | 2,529 | 50.2% | 1.6% |
| 35-49 Years | 1,061 | 88.7% | 1.0% | 1,061 | 84.9% | 1.1% | 3,171 | 75.1% | 1.0% | 5,887 | 57.2% | 1.1% |
| 50-64 Years | 344 | 87.8% | 1.8% | 344 | 83.3% | 2.0% | 909 | 72.8% | 1.9% | 1,403 | 54.1% | 2.9% |
| 65+ Years | 5 | + | + | 5 | + | + | 11 | 72.7% | 15.9% | 13 | 65.6% | 15.4% |
| Recipient Ethnicity/Race | | | | | | | | | | | | |
| White | 1,306 | 88.8% | 0.9% | 1,306 | 85.0% | 1.0% | 3,961 | 74.2% | 0.9% | 7,710 | 55.9% | 1.0% |
| African American | 265 | 88.1% | 2.0% | 265 | 83.1% | 2.3% | 780 | 66.7% | 2.4% | 1,256 | 50.1% | 2.6% |
| Hispanic/Latino | 148 | 90.5% | 2.4% | 148 | 86.4% | 2.8% | 452 | 76.6% | 2.5% | 706 | 57.6% | 3.8% |
| Asian | 23 | 91.3% | 5.7% | 23 | 91.3% | 5.7% | 75 | 84.0% | 4.5% | 108 | 59.2% | 11.7% |
| Other/Multi-race | 19 | 78.6% | 9.1% | 19 | 73.2% | 9.9% | 36 | 75.6% | 6.9% | 56 | 40.0% | 12.4% |
| Recipient Gender | | | | | | | | | | | | |
| Female | 650 | 89.0% | 1.2% | 650 | 84.9% | 1.4% | 2,012 | 70.6% | 1.3% | 3,899 | 54.1% | 1.4% |
| Male | 1,111 | 88.6% | 1.0% | 1,111 | 84.7% | 1.1% | 3,292 | 75.2% | 1.0% | 5,937 | 55.9% | 1.1% |

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney-pancreas and pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 8.8 (Continued)
Adjusted Pancreas Graft Survival, Kidney-Pancreas Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|--------------------------------|------------------|--------|-----------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2006 - 2007) | | | (Tx 2006 - 2007) | | | (Tx 2002 - 2007) | | | (Tx 1997 - 2007) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 1,761 | 88.7% | 0.8% | 1,761 | 84.8% | 0.9% | 5,304 | 73.5% | 0.8% | 9,836 | 55.3% | 0.9% |
| Transplant Center State | | | | | | | | | | | | |
| Alabama | 37 | 94.6% | 3.7% | 37 | 94.6% | 3.7% | 97 | 69.7% | 7.4% | 167 | 57.7% | 7.3% |
| Arkansas | 5 | 80.1% | 15.9% | 5 | 80.1% | 15.9% | 18 | 64.8% | 11.4% | 46 | 29.4% | 12.1% |
| Arizona | 70 | 90.0% | 3.6% | 70 | 87.1% | 4.0% | 132 | 74.3% | 5.6% | 182 | 60.6% | 7.0% |
| California | 214 | 88.8% | 2.1% | 214 | 85.0% | 2.4% | 660 | 77.3% | 2.0% | 1,199 | 59.0% | 2.8% |
| Colorado | 25 | 84.0% | 7.2% | 25 | 80.0% | 7.8% | 75 | 83.1% | 4.8% | 154 | 56.6% | 7.1% |
| Connecticut | 2 | 49.9% | 25.6% | 2 | 49.9% | 25.6% | 9 | 66.9% | 14.5% | 25 | 32.4% | 15.5% |
| District of Columbia | 13 | 92.3% | 7.1% | 13 | 76.9% | 11.2% | 51 | 67.5% | 8.9% | 119 | 49.2% | 7.0% |
| Florida | 133 | 91.7% | 2.4% | 133 | 87.7% | 2.9% | 428 | 74.3% | 2.7% | 648 | 56.8% | 4.1% |
| Georgia | 48 | 85.3% | 5.1% | 48 | 83.2% | 5.4% | 150 | 75.6% | 4.2% | 274 | 56.5% | 4.9% |
| Hawaii | 5 | + | + | 5 | + | + | 13 | 72.8% | 13.3% | 20 | 0.0% | 0.0% |
| Iowa | 13 | 77.1% | 11.1% | 13 | 77.1% | 11.1% | 53 | 84.9% | 4.9% | 118 | 71.1% | 5.1% |
| Illinois | 77 | 92.2% | 3.0% | 77 | 89.6% | 3.5% | 350 | 72.7% | 2.9% | 703 | 54.3% | 3.2% |
| Indiana | 59 | 89.8% | 3.9% | 59 | 89.8% | 3.9% | 151 | 76.3% | 4.9% | 183 | 66.9% | 7.0% |
| Kansas | 15 | 86.6% | 8.5% | 15 | 78.7% | 10.5% | 45 | 87.2% | 5.4% | 94 | 56.5% | 10.9% |
| Kentucky | 25 | 92.0% | 5.3% | 25 | 88.0% | 6.4% | 82 | 62.5% | 6.7% | 161 | 23.7% | 8.5% |
| Louisiana | 41 | 92.7% | 4.0% | 41 | 90.3% | 4.6% | 149 | 76.7% | 4.2% | 245 | 41.1% | 8.8% |
| Massachusetts | 7 | 85.8% | 12.2% | 7 | 85.8% | 12.2% | 22 | 81.8% | 8.0% | 47 | 59.9% | 10.7% |
| Maryland | 64 | 90.4% | 3.7% | 64 | 81.8% | 4.9% | 120 | 77.8% | 4.4% | 240 | 49.1% | 6.0% |
| Maine | 4 | + | + | 4 | + | + | 13 | 63.8% | 14.2% | 13 | 64.1% | 14.2% |
| Michigan | 51 | 74.5% | 6.0% | 51 | 70.5% | 6.3% | 110 | 66.8% | 6.5% | 173 | 60.0% | 4.5% |
| Minnesota | 45 | 78.0% | 6.1% | 45 | 73.6% | 6.5% | 132 | 65.5% | 4.7% | 354 | 48.0% | 3.4% |
| Missouri | 27 | 100.0% | 0.0% | 27 | 92.6% | 4.9% | 70 | 58.5% | 10.8% | 111 | 56.0% | 6.7% |
| North Carolina | 87 | 89.7% | 3.2% | 87 | 85.1% | 3.8% | 219 | 77.2% | 3.6% | 359 | 60.2% | 4.3% |
| North Dakota | 5 | 60.6% | 16.9% | 5 | 60.6% | 16.9% | 16 | 56.1% | 13.4% | 16 | 55.6% | 13.5% |
| Nebraska | 19 | + | + | 19 | + | + | 56 | 79.7% | 6.0% | 113 | 67.6% | 7.3% |
| New Hampshire | 7 | + | + | 7 | + | + | 7 | + | + | 7 | + | + |
| New Jersey | 42 | 88.1% | 4.9% | 42 | 88.1% | 4.9% | 99 | 66.6% | 6.6% | 180 | 58.4% | 6.0% |
| Nevada | 0 | - | - | 0 | - | - | 3 | + | + | 14 | 83.8% | 10.1% |
| New York | 34 | 94.1% | 4.0% | 34 | 85.3% | 6.0% | 110 | 65.5% | 6.1% | 262 | 55.7% | 4.4% |
| Ohio | 123 | 90.3% | 2.7% | 123 | 84.5% | 3.2% | 344 | 72.1% | 3.0% | 646 | 55.2% | 2.9% |
| Oklahoma | 12 | + | + | 12 | + | + | 34 | 73.5% | 7.4% | 56 | 62.1% | 11.5% |
| Oregon | 21 | 95.2% | 4.5% | 21 | 90.5% | 6.3% | 54 | 76.1% | 7.0% | 84 | 78.3% | 5.7% |
| Pennsylvania | 78 | 84.6% | 4.1% | 78 | 82.0% | 4.3% | 299 | 69.6% | 3.1% | 662 | 50.0% | 3.1% |

(Continued)

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney-pancreas and pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 8.8 (Continued)
Adjusted Pancreas Graft Survival, Kidney-Pancreas Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|--------------------------------|------------------|--------|-----------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2006 - 2007) | | | (Tx 2006 - 2007) | | | (Tx 2002 - 2007) | | | (Tx 1997 - 2007) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Transplant Center State | | | | | | | | | | | | |
| Puerto Rico | 5 | 80.1% | 15.9% | 5 | 80.1% | 15.9% | 5 | 78.9% | 16.8% | 5 | 79.5% | 16.4% |
| Rhode Island | 2 | 100.0% | 0.0% | 2 | 50.3% | 24.8% | 5 | 60.5% | 19.2% | 5 | 59.6% | 19.4% |
| South Carolina | 22 | 86.3% | 7.1% | 22 | 81.7% | 8.0% | 68 | 75.1% | 6.3% | 141 | 49.7% | 6.3% |
| South Dakota | 0 | - | - | 0 | - | - | 2 | + | + | 2 | + | + |
| Tennessee | 52 | 94.2% | 3.2% | 52 | 88.4% | 4.4% | 130 | 69.0% | 5.5% | 264 | 50.3% | 4.3% |
| Texas | 86 | 87.1% | 3.6% | 86 | 85.9% | 3.7% | 285 | 73.5% | 3.4% | 519 | 55.1% | 4.6% |
| Utah | 23 | 82.6% | 7.7% | 23 | 82.6% | 7.7% | 67 | 84.4% | 5.0% | 104 | 72.0% | 6.2% |
| Virginia | 28 | 82.1% | 7.1% | 28 | 71.3% | 8.4% | 77 | 64.8% | 6.3% | 164 | 51.4% | 6.4% |
| Vermont | 5 | + | + | 5 | + | + | 14 | 85.7% | 9.0% | 15 | 80.0% | 9.9% |
| Washington | 45 | 91.1% | 4.2% | 45 | 85.6% | 5.4% | 134 | 81.6% | 4.0% | 252 | 56.9% | 6.7% |
| Wisconsin | 85 | 83.6% | 4.0% | 85 | 77.6% | 4.5% | 346 | 72.6% | 2.8% | 690 | 57.3% | 2.9% |

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney-pancreas and pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.