

**Table 10.6b**  
**Immunosuppression Use for Induction by Discharge Regimen, 2004 to 2008**  
**Recipients with Intestine Transplants**

	Discharge Regimen (w/ or w/o Steroids Use)					All
	Tac	Tac+ MMF	Tac+ Aza	Tac+ Siro	Other	
<b>Functioning Graft with Immunosuppression Info at Discharge</b>	535	127	4	74	20	760
<b>% of All Regimens</b>	70.4%	16.7%	0.5%	9.7%	2.6%	100.0%
<b>Induction Drug</b>						
Atgam/NRATG/NRATS	0.2%	2.4%	0.0%	1.4%	20.0%	1.2%
OKT3	3.6%	40.9%	0.0%	8.1%	85.0%	12.4%
Thymoglobulin	10.1%	60.6%	0.0%	28.4%	205.0%	25.4%
Zenapax	10.1%	19.7%	0.0%	8.1%	100.0%	13.8%
Simulect	2.1%	0.8%	0.0%	0.0%	20.0%	2.1%
Campath	25.2%	1.6%	25.0%	6.8%	90.0%	21.2%
<b>No Induction Drugs Recorded</b>	52.5%	16.5%	75.0%	55.4%	65.0%	47.2%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.