

Table 11.6b
Immunosuppression Use for Induction by Discharge Regimen, 2004 to 2008
Recipients with Heart Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	91	266	2,955	5,306	221	118	78	8	76	754	9,873
% of All Regimens	0.9%	2.7%	29.9%	53.7%	2.2%	1.2%	0.8%	0.1%	0.8%	7.6%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	8.8%	7.1%	5.3%	4.9%	9.5%	6.8%	7.7%	37.5%	5.3%	16.6%	6.2%
OKT3	2.2%	0.0%	2.6%	1.8%	28.1%	5.9%	1.3%	0.0%	3.9%	6.9%	3.0%
Thymoglobulin	9.9%	11.3%	14.0%	18.8%	20.4%	38.1%	21.8%	12.5%	55.3%	25.2%	18.1%
Zenapax	4.4%	1.1%	21.1%	10.6%	3.2%	2.5%	9.0%	12.5%	0.0%	31.0%	14.7%
Simulect	13.2%	8.6%	13.1%	14.9%	5.0%	3.4%	12.8%	25.0%	1.3%	21.6%	14.2%
Campath	1.1%	7.9%	0.1%	3.0%	0.5%	0.0%	6.4%	0.0%	0.0%	2.9%	2.2%
No Induction Drugs Recorded	64.8%	65.0%	45.7%	49.7%	34.4%	43.2%	43.6%	37.5%	35.5%	47.2%	48.3%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.