

**Table 11.6d**  
**Immunosuppression Use for Maintenance by Regimen**  
**Prior to Discharge, 1999 to 2008**  
**Recipients with Heart Transplants**

	Transplant Year									
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
<b>Functioning Graft at Discharge</b>	1,922	1,982	2,008	1,977	1,899	1,867	1,958	2,040	2,070	2,039
<b>With Immunosuppression Info</b>	1,912	1,976	1,983	1,966	1,885	1,853	1,936	1,999	2,059	2,026
CyA	0.5%	0.7%	0.3%	0.1%	0.1%	0.1%	0.1%	0.2%	0.0%	0.0%
Tac	0.1%	0.1%	0.3%	0.1%	0.4%	0.2%	0.4%	0.6%	0.6%	1.1%
--- + MMF/MPA	0.1%	0.1%	0.0%	0.0%	0.1%	0.1%	0.1%	0.2%	0.0%	0.2%
CyA + MMF/MPA	1.0%	2.1%	0.9%	0.8%	2.0%	1.9%	1.6%	1.9%	2.4%	1.9%
Tac + MMF/MPA	0.1%	0.3%	0.4%	0.9%	2.7%	2.6%	2.7%	3.8%	7.0%	8.4%
--- + OtherAntimet	0.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%
CyA + OtherAntimet	0.8%	1.2%	0.5%	0.6%	0.5%	0.8%	0.4%	0.2%	0.1%	0.2%
Tac + OtherAntimet	0.2%	0.1%	0.1%	0.1%	0.1%	0.2%	0.2%	0.1%	0.2%	0.1%
--- + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%
CyA + MMF/MPA + OtherAntimet	0.5%	0.1%	0.0%	0.1%	0.9%	1.1%	0.3%	0.1%	0.0%	0.0%
Tac + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.0%	0.1%	0.0%	0.0%
CyA + Siro/Evero	0.0%	0.1%	0.0%	0.1%	0.1%	0.0%	0.1%	0.1%	0.1%	0.0%
Tac + Siro/Evero	0.0%	0.0%	0.1%	0.2%	0.1%	0.1%	0.3%	0.4%	0.2%	0.2%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.1%	0.1%	0.0%	0.0%
CyA + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.0%	0.0%	0.1%	0.0%
Tac + MMF/MPA + Siro/Evero	0.0%	0.1%	0.1%	0.1%	0.1%	0.2%	0.2%	0.1%	0.0%	0.0%
--- + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%
--- + Steroids	1.6%	2.5%	0.8%	1.9%	1.1%	0.6%	0.4%	0.5%	0.4%	0.5%
CyA + Steroids	10.8%	3.8%	1.5%	2.3%	1.8%	0.6%	0.7%	1.3%	0.9%	0.6%
Tac + Steroids	2.2%	1.9%	3.4%	3.7%	2.8%	2.1%	2.0%	1.9%	2.3%	2.3%
--- + MMF/MPA + Steroids	0.8%	1.1%	0.4%	0.8%	0.4%	0.8%	1.1%	0.9%	1.0%	0.8%
CyA + MMF/MPA + Steroids	39.7%	53.7%	51.3%	42.0%	41.7%	38.8%	31.1%	28.5%	24.1%	18.6%
Tac + MMF/MPA + Steroids	8.1%	8.7%	19.3%	23.7%	27.5%	36.3%	46.0%	48.8%	53.0%	58.4%
--- + OtherAntimet + Steroids	0.6%	0.2%	0.1%	0.1%	0.0%	0.5%	0.8%	0.1%	0.1%	0.0%
CyA + OtherAntimet + Steroids	24.5%	14.6%	10.0%	6.6%	4.4%	3.8%	2.1%	2.0%	1.3%	0.6%
Tac + OtherAntimet + Steroids	1.9%	2.1%	1.4%	1.4%	1.4%	1.0%	0.9%	1.3%	0.8%	1.3%
--- + Siro/Evero + Steroids	0.0%	0.2%	0.2%	0.2%	0.1%	0.1%	0.1%	0.1%	0.1%	0.0%
CyA + Siro/Evero + Steroids	1.0%	0.9%	0.2%	0.9%	2.0%	0.3%	0.4%	2.5%	2.8%	2.4%
Tac + Siro/Evero + Steroids	0.1%	0.9%	2.1%	5.5%	1.9%	0.9%	0.8%	0.9%	0.1%	0.0%
Other Regimen	5.0%	4.8%	6.9%	8.1%	7.9%	6.6%	7.2%	4.0%	1.9%	2.0%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Percentages in each column sum to 100 percent.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.