

**Table 11.6f**  
**Immunosuppression Use for Maintenance by Regimen**  
**One Year Following Transplantation, 1998 to 2007**  
**Recipients with Heart Transplants**

	Transplant Year									
	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
<b>Transplants</b>	2,348	2,188	2,199	2,202	2,155	2,057	2,015	2,125	2,193	2,209
<b>Functioning Graft at 1 Year PostTx</b>	1,940	1,780	1,820	1,850	1,823	1,755	1,748	1,807	1,899	1,924
<b>With Maintenance Use Recorded at 1 Year PostTx</b>	1,084	1,724	1,755	1,829	1,787	1,718	1,739	1,795	1,880	1,896
CyA	1.0%	0.9%	1.3%	0.8%	0.8%	0.8%	0.3%	0.3%	0.3%	0.4%
Tac	1.2%	1.3%	1.1%	0.7%	1.6%	1.3%	1.2%	2.4%	2.8%	2.6%
--- + MMF/MPA	0.2%	0.1%	0.2%	0.1%	0.2%	0.4%	0.2%	0.1%	0.2%	0.2%
CyA + MMF/MPA	7.6%	8.5%	9.7%	9.0%	8.6%	8.5%	8.3%	7.6%	6.7%	5.3%
Tac + MMF/MPA	2.2%	2.8%	3.4%	6.9%	6.3%	10.4%	10.8%	15.4%	16.6%	20.0%
--- + OtherAntimet	0.2%	0.2%	0.2%	0.1%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet	3.7%	3.3%	2.3%	1.6%	2.4%	1.2%	1.4%	0.7%	1.1%	0.8%
Tac + OtherAntimet	0.5%	0.9%	0.9%	0.9%	1.1%	1.0%	1.3%	1.1%	1.2%	2.0%
--- + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.1%	0.0%
CyA + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.1%	0.1%	0.0%	0.1%	0.1%	0.0%	0.0%
Tac + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.0%	0.0%
--- + Siro/Evero	0.0%	0.0%	0.0%	0.1%	0.0%	0.2%	0.0%	0.0%	0.0%	0.1%
CyA + Siro/Evero	0.0%	0.1%	0.1%	0.2%	0.8%	0.8%	0.5%	0.2%	0.4%	0.4%
Tac + Siro/Evero	0.0%	0.0%	0.2%	0.2%	1.3%	1.5%	0.9%	1.5%	1.2%	1.1%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.1%	0.1%	0.2%	0.2%	0.7%	0.5%	0.4%
CyA + MMF/MPA + Siro/Evero	0.0%	0.1%	0.1%	0.1%	0.2%	0.5%	0.1%	0.2%	0.1%	0.2%
Tac + MMF/MPA + Siro/Evero	0.0%	0.1%	0.0%	0.1%	0.3%	0.2%	0.3%	0.3%	0.5%	0.5%
--- + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.1%	0.0%	0.1%
CyA + OtherAntimet + Siro/Evero	0.0%	0.1%	0.0%	0.0%	0.1%	0.0%	0.1%	0.0%	0.1%	0.0%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.1%	0.0%	0.1%	0.0%	0.1%	0.0%	0.1%	0.0%
--- + Steroids	5.4%	2.0%	2.2%	3.4%	0.4%	0.2%	0.4%	0.1%	0.2%	0.1%
CyA + Steroids	2.6%	4.9%	3.6%	1.9%	1.9%	1.9%	0.9%	1.4%	1.1%	1.0%
Tac + Steroids	2.4%	2.3%	2.3%	4.3%	3.5%	3.1%	2.6%	2.8%	2.7%	2.4%
--- + MMF/MPA + Steroids	0.6%	0.5%	0.8%	0.5%	0.6%	0.3%	0.6%	0.4%	0.3%	0.3%
CyA + MMF/MPA + Steroids	36.3%	35.1%	41.0%	33.5%	27.8%	24.9%	23.6%	20.7%	18.7%	16.1%
Tac + MMF/MPA + Steroids	13.8%	14.6%	16.9%	21.2%	24.9%	25.7%	32.1%	33.5%	35.4%	36.6%
--- + OtherAntimet + Steroids	0.1%	0.1%	0.1%	0.2%	0.1%	0.1%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet + Steroids	15.0%	11.3%	6.6%	4.6%	3.2%	2.0%	2.8%	0.9%	1.1%	0.6%
Tac + OtherAntimet + Steroids	4.2%	3.4%	1.9%	2.6%	1.9%	2.0%	1.6%	1.3%	1.5%	1.6%
--- + Siro/Evero + Steroids	0.0%	0.1%	0.2%	0.4%	0.3%	0.2%	0.1%	0.2%	0.2%	0.3%
CyA + Siro/Evero + Steroids	0.2%	1.2%	1.3%	0.8%	1.2%	2.5%	1.0%	0.6%	2.1%	2.4%
Tac + Siro/Evero + Steroids	0.3%	0.5%	1.0%	2.8%	4.5%	4.4%	2.2%	2.1%	1.5%	0.9%
Other Regimen	2.6%	5.9%	2.7%	3.1%	6.0%	5.5%	6.3%	5.1%	3.5%	3.8%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.