

Table 12.6b
Immunosuppression Use for Induction by Discharge Regimen, 2004 to 2008
Recipients with Lung Transplants

	Discharge Regimen (w/ or w/o Steroids Use)									All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	71	359	368	3,142	572	1,605	9	21	214	6,361
% of All Regimens	1.1%	5.6%	5.8%	49.4%	9.0%	25.2%	0.1%	0.3%	3.4%	100.0%
Induction Drug										
Atgam/NRATG/NRATS	2.8%	4.7%	1.9%	6.4%	4.2%	6.4%	0.0%	4.8%	11.7%	6.0%
OKT3	1.4%	0.0%	2.2%	0.2%	6.6%	0.2%	0.0%	0.0%	1.9%	1.0%
Thymoglobulin	1.4%	2.8%	3.3%	5.6%	5.6%	0.8%	11.1%	4.8%	8.4%	4.1%
Zenapax	0.0%	3.9%	12.0%	17.5%	6.5%	15.0%	22.2%	4.8%	47.2%	15.5%
Simulect	7.0%	6.7%	18.8%	22.7%	14.2%	36.0%	44.4%	61.9%	70.1%	25.7%
Campath	5.6%	20.6%	3.0%	10.6%	0.0%	0.1%	0.0%	4.8%	25.7%	7.6%
No Induction Drugs Recorded	81.7%	61.8%	59.8%	38.2%	63.8%	42.0%	22.2%	23.8%	48.1%	44.8%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.