

**Table 12.6f**  
**Immunosuppression Use for Maintenance by Regimen**  
**One Year Following Transplantation, 1998 to 2007**  
**Recipients with Lung Transplants**

	Transplant Year									
	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
<b>Transplants</b>	893	922	977	1,083	1,054	1,099	1,185	1,407	1,406	1,471
<b>Functioning Graft at 1 Year PostTx</b>	631	668	735	805	834	900	985	1,132	1,156	1,161
<b>With Maintenance Use Recorded at 1 Year PostTx</b>	325	621	695	767	810	868	976	1,118	1,142	1,150
CyA	0.0%	0.0%	0.4%	0.0%	0.1%	0.0%	0.0%	0.0%	0.2%	0.1%
Tac	0.3%	0.2%	0.0%	0.3%	0.6%	0.3%	0.1%	0.0%	0.3%	0.3%
--- + MMF/MPA	0.3%	0.3%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.0%
CyA + MMF/MPA	0.0%	0.0%	0.1%	0.4%	0.1%	0.3%	0.0%	0.1%	0.2%	0.0%
Tac + MMF/MPA	0.6%	0.3%	1.3%	0.5%	1.0%	0.5%	0.4%	0.5%	1.2%	0.6%
--- + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet	0.3%	0.2%	0.1%	0.1%	0.4%	0.2%	0.0%	0.0%	0.1%	0.0%
Tac + OtherAntimet	0.0%	0.2%	0.1%	0.1%	1.2%	0.5%	0.2%	0.1%	0.2%	0.2%
CyA + MMF/MPA + OtherAntimet	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%
--- + Siro/Evero	0.0%	0.3%	0.0%	0.3%	0.1%	0.1%	0.0%	0.0%	0.0%	0.0%
CyA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%
Tac + Siro/Evero	0.0%	0.2%	0.1%	0.1%	0.2%	0.1%	0.0%	0.1%	0.1%	0.0%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.1%	0.1%	0.0%	0.0%	0.1%	0.0%	0.1%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.1%	0.1%	0.1%
--- + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.2%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.0%	0.0%	0.0%
--- + Steroids	0.9%	0.5%	0.4%	0.5%	0.4%	0.7%	0.4%	0.4%	0.4%	0.3%
CyA + Steroids	4.3%	5.6%	3.5%	2.1%	1.6%	2.3%	1.1%	1.3%	1.5%	2.3%
Tac + Steroids	5.5%	6.4%	5.5%	9.9%	7.3%	8.2%	9.9%	9.6%	8.5%	13.3%
--- + MMF/MPA + Steroids	0.3%	0.0%	0.4%	0.3%	0.1%	0.3%	0.2%	0.2%	0.4%	0.3%
CyA + MMF/MPA + Steroids	15.4%	17.4%	14.0%	14.0%	10.7%	8.9%	7.6%	7.6%	4.1%	3.7%
Tac + MMF/MPA + Steroids	17.5%	17.2%	26.6%	29.7%	32.7%	34.1%	40.9%	45.9%	47.7%	48.6%
--- + OtherAntimet + Steroids	0.3%	0.2%	0.4%	0.3%	0.2%	0.3%	0.1%	0.1%	0.4%	0.2%
CyA + OtherAntimet + Steroids	32.3%	26.7%	22.9%	12.4%	8.1%	8.8%	6.1%	4.9%	4.6%	2.9%
Tac + OtherAntimet + Steroids	18.2%	17.4%	16.3%	17.3%	20.1%	22.8%	23.0%	20.4%	22.2%	18.5%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.1%	0.7%	0.2%	0.2%	0.4%	0.1%	0.1%
CyA + Siro/Evero + Steroids	0.3%	1.8%	1.3%	0.9%	0.9%	0.5%	0.3%	0.1%	0.4%	0.1%
Tac + Siro/Evero + Steroids	0.0%	3.2%	2.7%	3.5%	6.3%	4.0%	4.2%	3.8%	3.4%	2.8%
Other Regimen	3.4%	1.9%	3.5%	7.0%	6.5%	6.5%	5.1%	4.5%	3.8%	5.7%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.