

Table 13.6b
Immunosuppression Use for Induction by Discharge Regimen, 2004 to 2008
Recipients with Heart-Lung Transplants

	Discharge Regimen (w/ or w/o Steroids Use)								All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	2	11	13	75	19	11	1	4	136
% of All Regimens	1.5%	8.1%	9.6%	55.1%	14.0%	8.1%	0.7%	2.9%	100.0%
Induction Drug									
Atgam/NRATG/NRATS	0.0%	9.1%	15.4%	14.7%	31.6%	0.0%	0.0%	50.0%	16.2%
OKT3	0.0%	9.1%	7.7%	1.3%	5.3%	0.0%	0.0%	0.0%	2.9%
Thymoglobulin	0.0%	0.0%	0.0%	9.3%	0.0%	9.1%	100.0%	0.0%	6.6%
Zenapax	0.0%	18.2%	23.1%	17.3%	26.3%	18.2%	0.0%	75.0%	20.6%
Simulect	0.0%	9.1%	38.5%	10.7%	15.8%	36.4%	0.0%	50.0%	16.9%
Campath	0.0%	18.2%	0.0%	20.0%	0.0%	0.0%	0.0%	75.0%	14.7%
No Induction Drugs Recorded	100.0%	54.5%	23.1%	29.3%	26.3%	36.4%	0.0%	0.0%	30.9%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.