

Table 13.6d
Immunosuppression Use for Maintenance by Regimen
Prior to Discharge, 1999 to 2008
Recipients with Heart-Lung Transplants

	Transplant Year									
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Functioning Graft at Discharge	34	33	21	22	17	35	29	24	29	21
With Immunosuppression Info	33	33	21	22	17	35	29	24	28	20
Tac	0.0%	0.0%	0.0%	0.0%	5.9%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + Steroids	0.0%	0.0%	4.8%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + Steroids	12.1%	3.0%	0.0%	0.0%	0.0%	2.9%	0.0%	4.2%	0.0%	0.0%
Tac + Steroids	3.0%	3.0%	0.0%	4.5%	11.8%	11.4%	3.4%	4.2%	14.3%	5.0%
CyA + MMF/MPA + Steroids	30.3%	36.4%	33.3%	22.7%	17.6%	17.1%	10.3%	8.3%	3.6%	5.0%
Tac + MMF/MPA + Steroids	18.2%	9.1%	19.0%	18.2%	5.9%	34.3%	41.4%	62.5%	67.9%	85.0%
CyA + OtherAntimet + Steroids	18.2%	33.3%	38.1%	27.3%	23.5%	14.3%	34.5%	12.5%	0.0%	5.0%
Tac + OtherAntimet + Steroids	6.1%	6.1%	4.8%	18.2%	29.4%	14.3%	6.9%	4.2%	10.7%	0.0%
Tac + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	2.9%	0.0%	0.0%	0.0%	0.0%
Other Regimen	12.1%	9.1%	0.0%	9.1%	5.9%	2.9%	3.4%	4.2%	3.6%	0.0%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Percentages in each column sum to 100 percent.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.
 "Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.