

Table 6.6b
Immunosuppression Use for Induction by Discharge Regimen, 2004 to 2008
Recipients with Pancreas Transplant Alone (PTA) Transplants

	Discharge Regimen (w/ or w/o Steroids Use)									All
	CyA	Tac	CyA+ MMF	Tac+ MMF	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	1	278	6	398	4	2	2	61	135	887
% of All Regimens	0.1%	31.3%	0.7%	44.9%	0.5%	0.2%	0.2%	6.9%	15.2%	100.0%
Induction Drug										
Atgam/NRATG/NRATS	0.0%	0.0%	0.0%	2.3%	0.0%	0.0%	0.0%	0.0%	0.7%	1.1%
OKT3	0.0%	4.7%	0.0%	5.3%	0.0%	0.0%	0.0%	3.3%	4.4%	4.7%
Thymoglobulin	0.0%	10.4%	100.0%	61.6%	25.0%	50.0%	100.0%	16.4%	111.9%	50.2%
Zenapax	0.0%	15.1%	16.7%	4.5%	0.0%	0.0%	0.0%	4.9%	7.4%	8.3%
Simulect	100.0%	0.7%	0.0%	4.8%	0.0%	0.0%	0.0%	0.0%	2.2%	2.8%
Campath	0.0%	28.4%	16.7%	15.3%	50.0%	0.0%	0.0%	45.9%	45.2%	26.2%
No Induction Drugs Recorded	0.0%	45.3%	0.0%	14.8%	25.0%	50.0%	0.0%	36.1%	10.4%	25.1%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.