

Table 7.6b
Immunosuppression Use for Induction by Discharge Regimen, 2004 to 2008
Recipients with Pancreas After Kidney (PAK) Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	4	50	47	991	1	6	18	22	76	134	1,349
% of All Regimens	0.3%	3.7%	3.5%	73.5%	0.1%	0.4%	1.3%	1.6%	5.6%	9.9%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	0.0%	0.0%	0.0%	1.8%	0.0%	0.0%	5.6%	0.0%	1.3%	1.5%	1.6%
OKT3	0.0%	0.0%	0.0%	0.3%	0.0%	0.0%	0.0%	0.0%	1.3%	1.5%	0.4%
Thymoglobulin	100.0%	22.0%	72.3%	55.8%	100.0%	66.7%	66.7%	100.0%	42.1%	120.1%	61.8%
Zenapax	0.0%	4.0%	6.4%	4.7%	0.0%	0.0%	0.0%	0.0%	2.6%	12.7%	5.3%
Simulect	0.0%	4.0%	6.4%	5.7%	0.0%	0.0%	5.6%	4.5%	3.9%	8.2%	5.7%
Campath	25.0%	48.0%	14.9%	11.9%	0.0%	16.7%	22.2%	0.0%	26.3%	47.8%	17.7%
No Induction Drugs Recorded	0.0%	26.0%	10.6%	22.6%	0.0%	16.7%	16.7%	0.0%	25.0%	7.5%	20.4%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.