

Table 7.6g
Immunosuppression Use for Maintenance Between Discharge
and One Year Following Transplantation, 1998 to 2007
Recipients with Pancreas After Kidney (PAK) Transplants

	Year of Transplant									
	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
Transplants	156	220	305	305	375	344	420	343	293	261
Tx with Follow-up Immunosuppression Info	110	167	220	245	272	258	320	255	221	206
Corticosteroids										
Any in Category	100.0%	100.0%	98.6%	98.8%	94.5%	78.7%	73.8%	74.9%	71.0%	59.2%
Steroids	100.0%	100.0%	98.6%	98.8%	94.5%	78.7%	73.8%	74.9%	71.0%	59.2%
Cyclosporine										
Any in Category	32.7%	14.4%	16.4%	15.1%	15.4%	11.2%	8.1%	6.7%	6.8%	5.8%
Sandimmune	5.5%	2.4%	3.2%	0.8%	1.1%	0.0%	0.3%	0.0%	0.5%	0.5%
Neoral	27.3%	12.0%	12.3%	12.7%	12.1%	9.3%	5.0%	6.3%	6.3%	5.3%
Gengraf	0.0%	0.0%	0.9%	0.8%	2.2%	1.6%	2.8%	0.4%	0.0%	0.0%
Eon	0.0%	0.0%	0.0%	0.8%	0.0%	0.4%	0.0%	0.0%	0.0%	0.0%
Tacrolimus										
Any in Category	76.4%	91.0%	91.8%	91.8%	87.1%	82.9%	87.2%	92.5%	91.9%	93.7%
Tacrolimus	76.4%	91.0%	91.8%	91.8%	87.1%	82.9%	87.2%	92.5%	91.9%	92.7%
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.0%
Antimetabolites										
Any in Category	99.1%	95.2%	84.5%	80.4%	80.5%	84.9%	82.5%	90.6%	87.3%	90.3%
Mycophenolate Mofetil	89.1%	86.2%	78.6%	78.0%	78.3%	81.0%	79.4%	84.3%	75.6%	75.7%
Mycophenolate Sodium	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	2.2%	4.3%	8.6%	12.1%
Azathioprine	10.0%	9.0%	5.9%	2.4%	2.2%	3.5%	0.3%	2.0%	1.8%	1.0%
Leflunomide	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.6%	0.0%	1.4%	1.5%
Cytosan	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
mTOR Inhibitors										
Any in Category	0.9%	7.2%	24.5%	30.6%	31.6%	27.5%	25.6%	19.2%	17.2%	18.9%
Sirolimus	0.9%	7.2%	24.5%	30.6%	31.6%	27.5%	25.6%	19.2%	17.2%	18.9%
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Percentages are calculated based on the total number of patients with any immunosuppression use between discharge and one year following transplantation. Individual drug percentages will not necessarily add up to the drug category percentages, as patients may be prescribed more than one drug within the same category.