

Table 8.6b
Immunosuppression Use for Induction by Discharge Regimen, 2004 to 2008
Recipients with Kidney-Pancreas Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	1	153	72	3,335	1	12	53	149	219	272	4,267
% of All Regimens	0.0%	3.6%	1.7%	78.2%	0.0%	0.3%	1.2%	3.5%	5.1%	6.4%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	0.0%	2.6%	1.4%	1.9%	0.0%	0.0%	0.0%	0.0%	0.5%	1.8%	1.8%
OKT3	0.0%	1.3%	0.0%	0.7%	0.0%	0.0%	1.9%	0.0%	1.8%	2.9%	0.9%
Thymoglobulin	100.0%	30.1%	66.7%	52.6%	0.0%	75.0%	20.8%	89.3%	43.4%	85.7%	54.6%
Zenapax	0.0%	6.5%	1.4%	6.3%	0.0%	0.0%	1.9%	0.0%	2.7%	2.6%	5.5%
Simulect	0.0%	2.6%	15.3%	9.2%	0.0%	16.7%	0.0%	0.7%	1.8%	5.5%	8.0%
Campath	0.0%	28.8%	1.4%	11.3%	0.0%	0.0%	69.8%	0.7%	27.9%	23.9%	13.7%
No Induction Drugs Recorded	0.0%	31.4%	16.7%	20.6%	100.0%	8.3%	9.4%	10.1%	26.0%	8.8%	19.9%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.