

Table 8.6e
Immunosuppression Use for Maintenance
Prior to Discharge, 1999 to 2008
Recipients with Kidney-Pancreas Transplants

	Year of Transplant									
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Functioning Graft at Discharge	907	888	861	881	847	864	884	905	840	820
With Immunosuppression Info	907	886	861	881	844	851	879	891	831	815
Corticosteroids										
Any in Category	97.9%	97.1%	93.3%	89.6%	83.8%	75.2%	71.0%	66.2%	65.5%	70.1%
Prednisone	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Steroids	97.9%	97.0%	93.3%	89.6%	83.8%	75.2%	71.0%	66.2%	65.5%	70.1%
Cyclosporine										
Any in Category	23.3%	18.8%	13.7%	9.8%	8.9%	6.2%	6.7%	6.4%	4.5%	4.4%
Sandimmune	2.4%	1.9%	0.9%	0.8%	0.1%	0.1%	0.2%	0.2%	0.2%	0.5%
Neoral	20.7%	16.1%	10.8%	6.8%	7.2%	5.3%	5.2%	5.4%	3.7%	3.7%
Gengraf	0.1%	0.8%	2.0%	1.9%	1.5%	0.8%	1.3%	0.8%	0.5%	0.2%
Eon	0.0%	0.0%	0.0%	0.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tacrolimus										
Any in Category	75.3%	82.3%	86.6%	90.2%	83.9%	84.6%	88.9%	92.4%	94.3%	94.0%
Tacrolimus	75.3%	82.3%	86.6%	90.2%	83.9%	84.6%	88.9%	92.1%	93.9%	93.3%
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.5%	0.7%
Antimetabolites										
Any in Category	92.7%	84.9%	79.9%	82.1%	82.7%	83.7%	81.8%	85.1%	91.3%	91.2%
Mycophenolate Mofetil	89.4%	82.4%	78.9%	81.0%	81.6%	81.8%	74.6%	71.6%	73.4%	72.5%
Mycophenolate Sodium	0.0%	0.0%	0.0%	0.0%	0.0%	1.5%	6.9%	12.3%	17.2%	18.5%
Azathioprine	3.3%	2.5%	1.0%	1.0%	1.1%	0.4%	0.2%	1.1%	0.7%	0.1%
Cytoxin	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
mTOR Inhibitors										
Any in Category	1.2%	13.2%	21.4%	19.0%	19.3%	16.8%	15.4%	16.3%	10.3%	8.0%
Sirolimus	1.0%	13.0%	21.4%	19.0%	19.2%	16.8%	15.4%	16.3%	10.3%	8.0%
Everolimus	0.2%	0.2%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Individual drug percentages will not necessarily add up to the drug category percentages, as patients may be prescribed more than one drug within the same category.