

Table 9.6b
Immunosuppression Use for Induction by Discharge Regimen, 2004 to 2008
Recipients with Liver Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	271	8,231	1,196	17,333	90	123	219	165	283	1,328	29,239
% of All Regimens	0.9%	28.2%	4.1%	59.3%	0.3%	0.4%	0.7%	0.6%	1.0%	4.5%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	0.0%	1.0%	0.4%	0.5%	1.1%	0.0%	0.0%	0.0%	0.4%	1.3%	0.7%
OKT3	0.4%	0.7%	0.5%	2.6%	0.0%	0.8%	0.9%	0.0%	2.5%	3.2%	1.9%
Thymoglobulin	5.2%	6.9%	7.6%	9.1%	13.3%	26.8%	16.0%	3.6%	18.7%	22.1%	9.2%
Zenapax	10.7%	4.1%	11.5%	6.0%	0.0%	6.5%	5.9%	58.2%	4.6%	16.3%	6.5%
Simulect	8.1%	4.3%	16.9%	6.4%	2.2%	28.5%	12.8%	4.2%	4.6%	17.1%	6.8%
Campath	0.7%	3.4%	0.3%	0.6%	0.0%	1.6%	0.0%	0.0%	1.8%	2.8%	1.5%
No Induction Drugs Recorded	75.3%	80.3%	64.0%	77.5%	83.3%	35.8%	65.3%	35.8%	70.3%	76.7%	77.1%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.